

6 Appendix

6.1 Proof of Lemma 3.1

Sketch of proof: We use change of variable $\mathbf{W} = \mathbf{Z}\mathbf{Z}^\top$. The matrix \mathbf{W} can be decomposed as $\mathbf{Z}\mathbf{Z}^\top$ where \mathbf{Z} is $p \times K$, if and only if $\mathbf{W} \succeq 0$ and has rank at most K . The constraints $\|\mathbf{Z}\|_2 \leq \tau$, $\|Z_i\|_2 \leq 1$ and $\|\mathbf{Z}\|_F \leq \beta$ are equivalent with the constraints $\mathbf{W} \preceq \tau^2 I$, $\text{diag}(\mathbf{W}) \leq 1$ and $\text{tr}(\mathbf{W}) \leq \beta^2$, respectively. Also, note that the condition that the regularization parameters of all (i, j) pairs of variables are non-negative is satisfied implicitly. For the diagonal elements, we have the constraint $\text{diag}(\mathbf{W}) \leq 1$ explicitly. Without loss of generality, assume $i < j$. We know $\mathbf{W} \succeq 0$, therefore we achieve: $M = \begin{bmatrix} W_{ii} & W_{ij} \\ W_{ji} & W_{jj} \end{bmatrix} \succeq 0$. Having $W_{ii} \leq 1$ and $W_{jj} \leq 1$, we conclude $W_{ij} = W_{ji} \leq 1$.

6.2 Proof of Lemma 3.3

Sketch of proof: In the \mathbf{Z} -step, to estimate \mathbf{Z} given Θ using the BCD method, we solve the following problem based on Eq (2):

$$\underset{\mathbf{Z} \in \mathcal{D}}{\text{maximize}} \quad \text{tr}(\mathbf{Z}\mathbf{Z}^\top |\Theta|), \quad (14)$$

which is equivalent to:

$$\underset{\mathbf{Z} \in \mathcal{D}}{\text{maximize}} \quad \sum_{i,j} (\mathbf{Z}\mathbf{Z}^\top)_{ij} |\Theta_{ij}|, \quad (15)$$

where \mathcal{D} is defined based on the two constraints (a) and (b) described above. First, note that by solving Eq (14), we will have $\|Z_i\|_2 = 1$, for all variables i . Hence, because of the binary assumption, each row of Z will have exactly one element with value 1 and the other elements will be 0.

Then, Eq (15) leads to:

$$\underset{\mathbf{Z} \in \mathcal{D}}{\text{maximize}} \quad \sum_{(i,j) \in \mathcal{F}_Z} |\Theta_{ij}|,$$

which is equivalent to

$$\underset{\mathbf{Z} \in \mathcal{D}}{\text{minimize}} \quad \sum_{(i,j) \notin \mathcal{F}_Z} |\Theta_{ij}|, \quad (16)$$

where \mathcal{F}_Z is the set of edges within blocks B_1, \dots, B_K : $\mathcal{F}_Z = \cup_{k=1}^K \{(i, j) | i, j \in B_k\}$. None of the blocks B_k for all k would be empty due to the constraint (b). Therefore, Eq (16) is equivalent with the K -way graph-cut problem on the similarity matrix $|\Theta|$.

Therefore, in the special case with constraints (a) and (b), \mathbf{Z} -step is K -way graph-cut on $|\Theta|$. The GRAB algorithm can be viewed as an iterative BCD method that 1) uses graph-cut as a clustering method to find \mathbf{Z} based on $|\Theta|$ as a similarity matrix, and 2) learns a network structure of the GGM by solving graphical lasso problem to find Θ .