
Supplementary Material: Improved Dynamic Regret for Non-degenerate Functions

Lijun Zhang*, **Tianbao Yang†**, **Jinfeng Yi‡**, **Rong Jin§**, **Zhi-Hua Zhou***

*National Key Laboratory for Novel Software Technology, Nanjing University, Nanjing, China

†Department of Computer Science, The University of Iowa, Iowa City, USA

‡AI Foundations Lab, IBM Thomas J. Watson Research Center, Yorktown Heights, NY, USA

§Alibaba Group, Seattle, USA

zhanglj@lamda.nju.edu.cn, tianbao-yang@uiowa.edu, jinfengyi@tencent.com

jinrong.jr@alibaba-inc.com, zhouzh@lamda.nju.edu.cn

A Proof of Theorem 1

For the sake of completeness, we include the proof of Theorem 1, which was proved by Mokhtari et al. [2016]. We need the following property of gradient descent.

Lemma 1. *Assume that $f : \mathcal{X} \mapsto \mathbb{R}$ is λ -strongly convex and L -smooth, and $\mathbf{x}_* = \operatorname{argmin}_{\mathbf{x} \in \mathcal{X}} f(\mathbf{x})$. Let $\mathbf{v} = \Pi_{\mathcal{X}}(\mathbf{u} - \eta \nabla f(\mathbf{u}))$, where $\eta \leq 1/L$. We have*

$$\|\mathbf{v} - \mathbf{x}_*\| \leq \sqrt{1 - \frac{2\lambda}{1/\eta + \lambda}} \|\mathbf{u} - \mathbf{x}_*\|.$$

The constant in the above lemma is better than that in Proposition 2 of Mokhtari et al. [2016].

Since $\|\nabla f_t(\mathbf{x})\| \leq G$ for any $t \in [T]$ and any $\mathbf{x} \in \mathcal{X}$, we have

$$\sum_{t=1}^T f_t(\mathbf{x}_t) - f_t(\mathbf{x}_t^*) \leq G \sum_{t=1}^T \|\mathbf{x}_t - \mathbf{x}_t^*\|. \quad (13)$$

We now proceed to bound $\sum_{t=1}^T \|\mathbf{x}_t - \mathbf{x}_t^*\|$. By the triangle inequality, we have

$$\sum_{t=1}^T \|\mathbf{x}_t - \mathbf{x}_t^*\| \leq \|\mathbf{x}_1 - \mathbf{x}_1^*\| + \sum_{t=2}^T (\|\mathbf{x}_t - \mathbf{x}_{t-1}^*\| + \|\mathbf{x}_{t-1}^* - \mathbf{x}_t^*\|). \quad (14)$$

Since

$$\mathbf{x}_t = \Pi_{\mathcal{X}}(\mathbf{x}_{t-1} - \eta \nabla f_{t-1}(\mathbf{x}_{t-1}))$$

using Lemma 1, we have

$$\|\mathbf{x}_t - \mathbf{x}_{t-1}^*\| \leq \gamma \|\mathbf{x}_{t-1} - \mathbf{x}_{t-1}^*\|. \quad (15)$$

From (14) and (15), we have

$$\sum_{t=1}^T \|\mathbf{x}_t - \mathbf{x}_t^*\| \leq \|\mathbf{x}_1 - \mathbf{x}_1^*\| + \gamma \sum_{t=2}^T \|\mathbf{x}_{t-1} - \mathbf{x}_{t-1}^*\| + \mathcal{P}_T^* \leq \|\mathbf{x}_1 - \mathbf{x}_1^*\| + \gamma \sum_{t=1}^T \|\mathbf{x}_t - \mathbf{x}_t^*\| + \mathcal{P}_T^*$$

implying

$$\sum_{t=1}^T \|\mathbf{x}_t - \mathbf{x}_t^*\| \leq \frac{1}{1-\gamma} \mathcal{P}_T^* + \frac{1}{1-\gamma} \|\mathbf{x}_1 - \mathbf{x}_1^*\|. \quad (16)$$

We complete the proof by substituting (16) into (13).

B Proof of Lemma 1

We first introduce the following property of strongly convex functions [Hazan and Kale, 2011].

Lemma 2. *Assume that $f : \mathcal{X} \mapsto \mathbb{R}$ is λ -strongly convex, and $\mathbf{x}_* = \operatorname{argmin}_{\mathbf{x} \in \mathcal{X}} f(\mathbf{x})$. Then, we have*

$$f(\mathbf{x}) - f(\mathbf{x}_*) \geq \frac{\lambda}{2} \|\mathbf{x} - \mathbf{x}_*\|^2, \quad \forall \mathbf{x} \in \mathcal{X}. \quad (17)$$

From the updating rule, we have

$$\mathbf{v} = \operatorname{argmin}_{\mathbf{x} \in \mathcal{X}} f(\mathbf{u}) + \langle \nabla f(\mathbf{u}), \mathbf{x} - \mathbf{u} \rangle + \frac{1}{2\eta} \|\mathbf{x} - \mathbf{u}\|^2.$$

According to Lemma 2, we have

$$\begin{aligned} & f(\mathbf{u}) + \langle \nabla f(\mathbf{u}), \mathbf{v} - \mathbf{u} \rangle + \frac{1}{2\eta} \|\mathbf{v} - \mathbf{u}\|^2 \\ & \leq f(\mathbf{u}) + \langle \nabla f(\mathbf{u}), \mathbf{x}_* - \mathbf{u} \rangle + \frac{1}{2\eta} \|\mathbf{x}_* - \mathbf{u}\|^2 - \frac{1}{2\eta} \|\mathbf{v} - \mathbf{x}_*\|^2. \end{aligned} \quad (18)$$

Since $f(\mathbf{x})$ is λ -strongly convex, we have

$$f(\mathbf{u}) + \langle \nabla f(\mathbf{u}), \mathbf{x}_* - \mathbf{u} \rangle \leq f(\mathbf{x}_*) - \frac{\lambda}{2} \|\mathbf{x}_* - \mathbf{u}\|^2. \quad (19)$$

On the other hand, the smoothness assumption implies

$$f(\mathbf{v}) \leq f(\mathbf{u}) + \langle \nabla f(\mathbf{u}), \mathbf{v} - \mathbf{u} \rangle + \frac{L}{2} \|\mathbf{v} - \mathbf{u}\|^2 \leq f(\mathbf{u}) + \langle \nabla f(\mathbf{u}), \mathbf{v} - \mathbf{u} \rangle + \frac{1}{2\eta} \|\mathbf{v} - \mathbf{u}\|^2. \quad (20)$$

Combining (18), (19), and (20), we obtain

$$f(\mathbf{v}) \leq f(\mathbf{x}_*) - \frac{\lambda}{2} \|\mathbf{x}_* - \mathbf{u}\|^2 + \frac{1}{2\eta} \|\mathbf{x}_* - \mathbf{u}\|^2 - \frac{1}{2\eta} \|\mathbf{v} - \mathbf{x}_*\|^2. \quad (21)$$

Applying Lemma 2 again, we have

$$f(\mathbf{v}) - f(\mathbf{x}_*) \geq \frac{\lambda}{2} \|\mathbf{v} - \mathbf{x}_*\|^2. \quad (22)$$

We complete the proof by substituting (22) into (21) and rearranging.

C Proof of Theorem 2

Since $f_t(\cdot)$ is L -smooth, we have

$$f_t(\mathbf{x}_t) - f_t(\mathbf{x}_t^*) \leq \langle \nabla f_t(\mathbf{x}_t^*), \mathbf{x}_t - \mathbf{x}_t^* \rangle + \frac{L}{2} \|\mathbf{x}_t - \mathbf{x}_t^*\|^2 \leq \|\nabla f_t(\mathbf{x}_t^*)\| \|\mathbf{x}_t - \mathbf{x}_t^*\| + \frac{L}{2} \|\mathbf{x}_t - \mathbf{x}_t^*\|^2.$$

Combining with the fact

$$\|\nabla f_t(\mathbf{x}_t^*)\| \|\mathbf{x}_t - \mathbf{x}_t^*\| \leq \frac{1}{2\alpha} \|\nabla f_t(\mathbf{x}_t^*)\|^2 + \frac{\alpha}{2} \|\mathbf{x}_t - \mathbf{x}_t^*\|^2$$

for any $\alpha > 0$, we obtain

$$f_t(\mathbf{x}_t) - f_t(\mathbf{x}_t^*) \leq \frac{1}{2\alpha} \|\nabla f_t(\mathbf{x}_t^*)\|^2 + \frac{L + \alpha}{2} \|\mathbf{x}_t - \mathbf{x}_t^*\|^2.$$

Summing the above inequality over $t = 1, \dots, T$, we get

$$\sum_{t=1}^T f_t(\mathbf{x}_t) - f_t(\mathbf{x}_t^*) \leq \frac{1}{2\alpha} \sum_{t=1}^T \|\nabla f_t(\mathbf{x}_t^*)\|^2 + \frac{L + \alpha}{2} \sum_{t=1}^T \|\mathbf{x}_t - \mathbf{x}_t^*\|^2. \quad (23)$$

We now proceed to bound $\sum_{t=1}^T \|\mathbf{x}_t - \mathbf{x}_t^*\|^2$. We have

$$\sum_{t=1}^T \|\mathbf{x}_t - \mathbf{x}_t^*\|^2 \leq \|\mathbf{x}_1 - \mathbf{x}_1^*\|^2 + 2 \sum_{t=2}^T (\|\mathbf{x}_t - \mathbf{x}_{t-1}^*\|^2 + \|\mathbf{x}_{t-1}^* - \mathbf{x}_t^*\|^2). \quad (24)$$

Recall the updating rule

$$\mathbf{z}_{t-1}^{j+1} = \Pi_{\mathcal{X}} \left(\mathbf{z}_{t-1}^j - \eta \nabla f_{t-1}(\mathbf{z}_{t-1}^j) \right), j = 1, \dots, K.$$

From Lemma 1, we have

$$\|\mathbf{z}_{t-1}^{j+1} - \mathbf{x}_{t-1}^*\|^2 \leq \left(1 - \frac{2\lambda}{1/\eta + \lambda} \right) \|\mathbf{z}_{t-1}^j - \mathbf{x}_{t-1}^*\|^2$$

which implies

$$\|\mathbf{x}_t - \mathbf{x}_{t-1}^*\|^2 = \|\mathbf{z}_{t-1}^{K+1} - \mathbf{x}_{t-1}^*\|^2 \leq \left(1 - \frac{2\lambda}{1/\eta + \lambda} \right)^K \|\mathbf{x}_{t-1} - \mathbf{x}_{t-1}^*\|^2 \leq \frac{1}{4} \|\mathbf{x}_{t-1} - \mathbf{x}_{t-1}^*\|^2 \quad (25)$$

where we choose $K = \lceil \frac{1/\eta + \lambda}{2\lambda} \ln 4 \rceil$ such that

$$\left(1 - \frac{2\lambda}{1/\eta + \lambda} \right)^K \leq \exp \left(-\frac{2K\lambda}{1/\eta + \lambda} \right) \leq \frac{1}{4}.$$

From (24) and (25), we have

$$\begin{aligned} \sum_{t=1}^T \|\mathbf{x}_t - \mathbf{x}_t^*\|^2 &\leq \|\mathbf{x}_1 - \mathbf{x}_1^*\|^2 + \frac{1}{2} \sum_{t=2}^T \|\mathbf{x}_{t-1} - \mathbf{x}_{t-1}^*\|^2 + 2\mathcal{S}_T^* \\ &\leq \|\mathbf{x}_1 - \mathbf{x}_1^*\|^2 + \frac{1}{2} \sum_{t=1}^T \|\mathbf{x}_t - \mathbf{x}_t^*\|^2 + 2\mathcal{S}_T^* \end{aligned}$$

implying

$$\sum_{t=1}^T \|\mathbf{x}_t - \mathbf{x}_t^*\|^2 \leq 4\mathcal{S}_T^* + 2\|\mathbf{x}_1 - \mathbf{x}_1^*\|^2.$$

Substituting the above inequality into (23), we have

$$\sum_{t=1}^T f_t(\mathbf{x}_t) - f_t(\mathbf{x}_t^*) \leq \frac{1}{2\alpha} \sum_{t=1}^T \|\nabla f_t(\mathbf{x}_t^*)\|^2 + 2(L + \alpha)\mathcal{S}_T^* + (L + \alpha)\|\mathbf{x}_1 - \mathbf{x}_1^*\|^2$$

for all $\alpha \geq 0$. Finally, we show that the dynamic regret can still be upper bounded by \mathcal{P}_T^* . From the previous analysis, we have

$$\|\mathbf{x}_t - \mathbf{x}_{t-1}^*\|^2 \stackrel{(25)}{\leq} \frac{1}{4} \|\mathbf{x}_{t-1} - \mathbf{x}_{t-1}^*\|^2 \Rightarrow \|\mathbf{x}_t - \mathbf{x}_{t-1}^*\| \leq \frac{1}{2} \|\mathbf{x}_{t-1} - \mathbf{x}_{t-1}^*\|.$$

Then, we can set $\gamma = 1/2$ in Theorem 1 and obtain

$$\sum_{t=1}^T f_t(\mathbf{x}_t) - f_t(\mathbf{x}_t^*) \leq 2G\mathcal{P}_T^* + 2G\|\mathbf{x}_1 - \mathbf{x}_1^*\|.$$

D Proof of Theorem 5

We will randomly generate a sequence of functions $f_t : \mathbb{R}^d \mapsto \mathbb{R}, t = 1, \dots, T$, where each $f_t(\cdot)$ is independently sampled from a distribution \mathcal{P} . For any deterministic algorithm \mathcal{A} , it generates a sequence of solutions $\mathbf{x}_t \in \mathcal{X}, t = 1, \dots, T$, we define the expected dynamic regret as

$$\mathbb{E}[R_T^*] = \mathbb{E} \left[\sum_{t=1}^T f_t(\mathbf{x}_t) - f_t(\mathbf{x}_t^*) \right].$$

We will show that there exists a distribution of strongly convex and smooth functions such that for any fixed algorithm \mathcal{A} , we have $\mathbb{E}[R_T^*] \geq \mathbb{E}[\mathcal{S}_T^*]$.

For each round t , we randomly sample a vector $\varepsilon_t \in \mathbb{R}^d$ from the Gaussian distribution $\mathcal{N}(0, I)$. Using ε_t , we create a function

$$f_t(\mathbf{x}) = 2 \|\mathbf{x} - \tau \varepsilon_t\|^2$$

which is both strongly convex and smooth. Notice that \mathbf{x}_t is independent from ε_t , and thus we can bound the expected dynamic regret as follows:

$$\mathbb{E}[R_T^*] = \sum_{t=1}^T \mathbb{E}[f_t(\mathbf{x}_t) - f_t(\mathbf{x}_t^*)] = 2 \sum_{t=1}^T \mathbb{E}[\|\mathbf{x}_t\|^2 + d\tau^2] \geq 2dT\tau^2.$$

We furthermore bound \mathcal{S}_T^* as follows

$$\mathbb{E}[\mathcal{S}_T^*] = \sum_{t=2}^T \mathbb{E}[\|\varepsilon_t - \varepsilon_{t-1}\|^2 \tau^2] = 2d(T-1)\tau^2.$$

Therefore, $\mathbb{E}[R_T^*] \geq \mathbb{E}[\mathcal{S}_T^*]$. Hence, for any given algorithm \mathcal{A} , there exists a sequence of functions f_1, \dots, f_T , such that $\sum_{t=1}^T f_t(\mathbf{x}_t) - f_t(\mathbf{x}_t^*) = \Omega(\mathcal{S}_T^*)$.

E Proof of Theorem 6

The proof is similar to that of Theorem 1.

We need the following property of gradient descent when applied to semi-strongly convex and smooth functions [Necoara et al., 2015], which is analogous to Lemma 1 developed for strongly convex functions.

Lemma 3. *Assume that $f(\cdot)$ is L -smooth and satisfies the semi-strong convexity condition in (8). Let $\mathbf{v} = \Pi_{\mathcal{X}}(\mathbf{u} - \eta \nabla f(\mathbf{u}))$, where $\eta \leq 1/L$. We have*

$$\|\mathbf{v} - \Pi_{\mathcal{X}^*}(\mathbf{v})\| \leq \sqrt{1 - \frac{\beta}{1/\eta + \beta}} \|\mathbf{u} - \Pi_{\mathcal{X}^*}(\mathbf{u})\|.$$

Since $\|\nabla f_t(\mathbf{x})\| \leq G$ for any $t \in [T]$ and any $\mathbf{x} \in \mathcal{X}$, we have

$$\sum_{t=1}^T f_t(\mathbf{x}_t) - \sum_{t=1}^T \min_{\mathbf{x} \in \mathcal{X}} f_t(\mathbf{x}) = \sum_{t=1}^T f_t(\mathbf{x}_t) - f_t(\Pi_{\mathcal{X}_t^*}(\mathbf{x}_t)) \leq G \sum_{t=1}^T \|\mathbf{x}_t - \Pi_{\mathcal{X}_t^*}(\mathbf{x}_t)\|. \quad (26)$$

We now proceed to bound $\sum_{t=1}^T \|\mathbf{x}_t - \Pi_{\mathcal{X}_t^*}(\mathbf{x}_t)\|$. By the triangle inequality, we have

$$\sum_{t=1}^T \|\mathbf{x}_t - \Pi_{\mathcal{X}_t^*}(\mathbf{x}_t)\| \leq \|\mathbf{x}_1 - \Pi_{\mathcal{X}_1^*}(\mathbf{x}_1)\| + \sum_{t=2}^T \left(\|\mathbf{x}_t - \Pi_{\mathcal{X}_{t-1}^*}(\mathbf{x}_t)\| + \|\Pi_{\mathcal{X}_{t-1}^*}(\mathbf{x}_t) - \Pi_{\mathcal{X}_t^*}(\mathbf{x}_t)\| \right). \quad (27)$$

Since

$$\mathbf{x}_t = \Pi_{\mathcal{X}}(\mathbf{x}_{t-1} - \eta \nabla f_{t-1}(\mathbf{x}_{t-1}))$$

using Lemma 3, we have

$$\|\mathbf{x}_t - \Pi_{\mathcal{X}_{t-1}^*}(\mathbf{x}_t)\| \leq \gamma \|\mathbf{x}_{t-1} - \Pi_{\mathcal{X}_{t-1}^*}(\mathbf{x}_{t-1})\|. \quad (28)$$

From (27) and (28), we have

$$\begin{aligned} & \sum_{t=1}^T \|\mathbf{x}_t - \Pi_{\mathcal{X}_t^*}(\mathbf{x}_t)\| \\ & \leq \|\mathbf{x}_1 - \Pi_{\mathcal{X}_1^*}(\mathbf{x}_1)\| + \gamma \sum_{t=2}^T \|\mathbf{x}_{t-1} - \Pi_{\mathcal{X}_{t-1}^*}(\mathbf{x}_{t-1})\| + \sum_{t=2}^T \|\Pi_{\mathcal{X}_{t-1}^*}(\mathbf{x}_t) - \Pi_{\mathcal{X}_t^*}(\mathbf{x}_t)\| \\ & \leq \|\mathbf{x}_1 - \Pi_{\mathcal{X}_1^*}(\mathbf{x}_1)\| + \gamma \sum_{t=1}^T \|\mathbf{x}_t - \Pi_{\mathcal{X}_t^*}(\mathbf{x}_t)\| + \mathcal{P}_T^* \end{aligned}$$

implying

$$\sum_{t=1}^T \|\mathbf{x}_t - \Pi_{\mathcal{X}_t^*}(\mathbf{x}_t)\| \leq \frac{1}{1-\gamma} \mathcal{P}_T^* + \frac{1}{1-\gamma} \|\mathbf{x}_1 - \Pi_{\mathcal{X}_1^*}(\mathbf{x}_1)\|. \quad (29)$$

We complete the proof by substituting (29) into (26).

F Proof of Lemma 3

For the sake of completeness, we provide the proof of Lemma 3, which can also be found in the work of Necoara et al. [2015].

The analysis is similar to that of Lemma 1. Define

$$\bar{\mathbf{u}} = \Pi_{\mathcal{X}^*}(\mathbf{u}), \text{ and } \bar{\mathbf{v}} = \Pi_{\mathcal{X}^*}(\mathbf{v}).$$

From the optimality condition of \mathbf{v} , we have

$$\begin{aligned} & f(\mathbf{u}) + \langle \nabla f(\mathbf{u}), \mathbf{v} - \mathbf{u} \rangle + \frac{1}{2\eta} \|\mathbf{v} - \mathbf{u}\|^2 \\ & \leq f(\mathbf{u}) + \langle \nabla f(\mathbf{u}), \bar{\mathbf{u}} - \mathbf{u} \rangle + \frac{1}{2\eta} \|\bar{\mathbf{u}} - \mathbf{u}\|^2 - \frac{1}{2\eta} \|\mathbf{v} - \bar{\mathbf{u}}\|^2. \end{aligned} \quad (30)$$

From the convexity of $f(\mathbf{x})$, we have

$$f(\mathbf{u}) + \langle \nabla f(\mathbf{u}), \bar{\mathbf{u}} - \mathbf{u} \rangle \leq f(\bar{\mathbf{u}}). \quad (31)$$

Combining (30), (31), and (20), we obtain

$$f(\mathbf{v}) \leq f(\bar{\mathbf{u}}) + \frac{1}{2\eta} \|\bar{\mathbf{u}} - \mathbf{u}\|^2 - \frac{1}{2\eta} \|\mathbf{v} - \bar{\mathbf{u}}\|^2. \quad (32)$$

From the semi-strong convexity of $f(\cdot)$, we further have

$$f(\mathbf{v}) - f(\bar{\mathbf{u}}) \geq \frac{\beta}{2} \|\mathbf{v} - \bar{\mathbf{v}}\|^2.$$

Substituting the above inequality into (32), we have

$$\frac{1}{2\eta} \|\bar{\mathbf{u}} - \mathbf{u}\|^2 \geq \frac{1}{2\eta} \|\mathbf{v} - \bar{\mathbf{u}}\|^2 + \frac{\beta}{2} \|\mathbf{v} - \bar{\mathbf{v}}\|^2 \geq \left(\frac{1}{2\eta} + \frac{\beta}{2} \right) \|\mathbf{v} - \bar{\mathbf{v}}\|^2$$

which completes the proof.

G Proof of Theorem 7

The proof is similar to that of Theorem 2. In the following, we just provide the key differences.

Following the derivation of (23), we get

$$\begin{aligned} \sum_{t=1}^T f_t(\mathbf{x}_t) - \sum_{t=1}^T \min_{\mathbf{x} \in \mathcal{X}} f_t(\mathbf{x}) & \leq \frac{1}{2\alpha} \sum_{t=1}^T \|\nabla f_t(\Pi_{\mathcal{X}_t^*}(\mathbf{x}_t))\|^2 + \frac{L + \alpha}{2} \sum_{t=1}^T \|\mathbf{x}_t - \Pi_{\mathcal{X}_t^*}(\mathbf{x}_t)\|^2 \\ & \leq \frac{1}{2\alpha} G_T^* + \frac{L + \alpha}{2} \sum_{t=1}^T \|\mathbf{x}_t - \Pi_{\mathcal{X}_t^*}(\mathbf{x}_t)\|^2 \end{aligned} \quad (33)$$

for any $\alpha > 0$.

To bound $\sum_{t=1}^T \|\mathbf{x}_t - \Pi_{\mathcal{X}_t^*}(\mathbf{x}_t)\|^2$, we have

$$\sum_{t=1}^T \|\mathbf{x}_t - \Pi_{\mathcal{X}_t^*}(\mathbf{x}_t)\|^2 \leq \|\mathbf{x}_1 - \Pi_{\mathcal{X}_1^*}(\mathbf{x}_1)\|^2 + 2 \sum_{t=2}^T \left(\left\| \mathbf{x}_t - \Pi_{\mathcal{X}_{t-1}^*}(\mathbf{x}_t) \right\|^2 + \left\| \Pi_{\mathcal{X}_{t-1}^*}(\mathbf{x}_t) - \Pi_{\mathcal{X}_t^*}(\mathbf{x}_t) \right\|^2 \right). \quad (34)$$

From Lemma 3 and the updating rule

$$\mathbf{z}_{t-1}^{j+1} = \Pi_{\mathcal{X}} \left(\mathbf{z}_{t-1}^j - \eta \nabla f_{t-1}(\mathbf{z}_{t-1}^j) \right), \quad j = 1, \dots, K$$

we have

$$\left\| \mathbf{z}_{t-1}^{j+1} - \Pi_{\mathcal{X}_{t-1}^*}(\mathbf{z}_{t-1}^{j+1}) \right\|^2 \leq \left(1 - \frac{\beta}{1/\eta + \beta} \right) \left\| \mathbf{z}_{t-1}^j - \Pi_{\mathcal{X}_{t-1}^*}(\mathbf{z}_{t-1}^j) \right\|^2, \quad j = 1, \dots, K$$

which implies

$$\begin{aligned} \left\| \mathbf{x}_t - \Pi_{\mathcal{X}_{t-1}^*}(\mathbf{x}_t) \right\|^2 &= \left\| \mathbf{z}_{t-1}^{K+1} - \Pi_{\mathcal{X}_{t-1}^*}(\mathbf{z}_{t-1}^{K+1}) \right\|^2 \\ &\leq \left(1 - \frac{\beta}{1/\eta + \beta} \right)^K \left\| \mathbf{x}_{t-1} - \Pi_{\mathcal{X}_{t-1}^*}(\mathbf{x}_{t-1}) \right\|^2 \leq \frac{1}{4} \left\| \mathbf{x}_{t-1} - \Pi_{\mathcal{X}_{t-1}^*}(\mathbf{x}_{t-1}) \right\|^2 \end{aligned} \quad (35)$$

where we choose $K = \lceil \frac{1/\eta + \beta}{\beta} \ln 4 \rceil$ such that

$$\left(1 - \frac{\beta}{1/\eta + \beta} \right)^K \leq \exp\left(-\frac{K\beta}{1/\eta + \beta}\right) \leq \frac{1}{4}.$$

From (34) and (35), we have

$$\begin{aligned} \sum_{t=1}^T \left\| \mathbf{x}_t - \Pi_{\mathcal{X}_t^*}(\mathbf{x}_t) \right\|^2 &\leq \left\| \mathbf{x}_1 - \Pi_{\mathcal{X}_1^*}(\mathbf{x}_1) \right\|^2 + \frac{1}{2} \sum_{t=2}^T \left\| \mathbf{x}_{t-1} - \Pi_{\mathcal{X}_{t-1}^*}(\mathbf{x}_{t-1}) \right\|^2 + 2\mathcal{S}_T^* \\ &\leq \left\| \mathbf{x}_1 - \Pi_{\mathcal{X}_1^*}(\mathbf{x}_1) \right\|^2 + \frac{1}{2} \sum_{t=1}^T \left\| \mathbf{x}_t - \Pi_{\mathcal{X}_t^*}(\mathbf{x}_t) \right\|^2 + 2\mathcal{S}_T^* \end{aligned} \quad (36)$$

implying

$$\sum_{t=1}^T \left\| \mathbf{x}_t - \Pi_{\mathcal{X}_t^*}(\mathbf{x}_t) \right\|^2 \leq 4\mathcal{S}_T^* + 2 \left\| \mathbf{x}_1 - \Pi_{\mathcal{X}_1^*}(\mathbf{x}_1) \right\|^2.$$

Substituting the above inequality into (33), we have

$$\sum_{t=1}^T f_t(\mathbf{x}_t) - \sum_{t=1}^T \min_{\mathbf{x} \in \mathcal{X}} f_t(\mathbf{x}) \leq \frac{1}{2\alpha} G_T^* + 2(L + \alpha)\mathcal{S}_T^* + (L + \alpha) \left\| \mathbf{x}_1 - \Pi_{\mathcal{X}_1^*}(\mathbf{x}_1) \right\|^2, \quad \forall \alpha \geq 0.$$

Finally, we show that the dynamic regret can still be upper bounded by \mathcal{P}_T^* . From the previous analysis, we have

$$\left\| \mathbf{x}_t - \Pi_{\mathcal{X}_{t-1}^*}(\mathbf{x}_t) \right\| \stackrel{(35)}{\leq} \frac{1}{2} \left\| \mathbf{x}_{t-1} - \Pi_{\mathcal{X}_{t-1}^*}(\mathbf{x}_{t-1}) \right\|.$$

Then, we can set $\gamma = 1/2$ in Theorem 6 and obtain

$$\sum_{t=1}^T f_t(\mathbf{x}_t) - \sum_{t=1}^T \min_{\mathbf{x} \in \mathcal{X}} f_t(\mathbf{x}) \leq 2G\mathcal{P}_T^* + 2G \left\| \mathbf{x}_1 - \Pi_{\mathcal{X}_1^*}(\mathbf{x}_1) \right\|.$$

H Proof of Theorem 8

The inequality (12) follows directly from the result in Section 2.2.X.C of Nemirovski [2004]. To prove the rest of this theorem, we will use the following properties of self-concordant functions and the damped Newton method [Nemirovski, 2004].

Lemma 4. *Let $f(\mathbf{x})$ be a self-concordant function, and $\|\mathbf{h}\|_{\mathbf{x}} = \sqrt{\mathbf{h}^\top \nabla^2 f(\mathbf{x}) \mathbf{h}}$. Then, all points within the Dikin ellipsoid $W_{\mathbf{x}}$ centered at \mathbf{x} , defined as $W_{\mathbf{x}} = \{\mathbf{x}' : \|\mathbf{x}' - \mathbf{x}\|_{\mathbf{x}} \leq 1\}$, share similar second order structure. More specifically, for a given point \mathbf{x} and for any \mathbf{h} with $\|\mathbf{h}\|_{\mathbf{x}} \leq 1$, we have*

$$(1 - \|\mathbf{h}\|_{\mathbf{x}})^2 \nabla^2 f(\mathbf{x}) \preceq \nabla^2 f(\mathbf{x} + \mathbf{h}) \preceq \frac{\nabla^2 f(\mathbf{x})}{(1 - \|\mathbf{h}\|_{\mathbf{x}})^2}. \quad (37)$$

Define $\mathbf{x}^* = \operatorname{argmin}_{\mathbf{x}} f(\mathbf{x})$. Then, we have

$$\|\mathbf{x} - \mathbf{x}^*\|_{\mathbf{x}^*} \leq \frac{\lambda(\mathbf{x})}{1 - \lambda(\mathbf{x})} \quad (38)$$

where $\lambda(\mathbf{x}) = \sqrt{\mathbf{x}^\top [\nabla^2 f(\mathbf{x})]^{-1} \mathbf{x}}$.

Consider the the damped Newton method: $\mathbf{v} = \mathbf{u} - \frac{1}{1+\lambda(\mathbf{u})} [\nabla^2 f(\mathbf{u})]^{-1} \nabla f(\mathbf{u})$. Then, we have

$$\lambda(\mathbf{v}) \leq 2\lambda^2(\mathbf{u}). \quad (39)$$

We will also use the following inequality frequently

$$\begin{aligned}
& \|\mathbf{x}\|_t^2 = \mathbf{x}^\top \nabla^2 f_t(\mathbf{x}_t^*) \mathbf{x} \\
& = \mathbf{x}^\top [\nabla^2 f_{t-1}(\mathbf{x}_{t-1}^*)]^{-\frac{1}{2}} [\nabla^2 f_{t-1}(\mathbf{x}_{t-1}^*)]^{-\frac{1}{2}} \nabla^2 f_t(\mathbf{x}_t^*) [\nabla^2 f_{t-1}(\mathbf{x}_{t-1}^*)]^{-\frac{1}{2}} [\nabla^2 f_{t-1}(\mathbf{x}_{t-1}^*)]^{-\frac{1}{2}} \mathbf{x} \\
& \stackrel{(10)}{\leq} \mu \mathbf{x}^\top \nabla^2 f_{t-1}(\mathbf{x}_{t-1}^*) \mathbf{x} = \mu \|\mathbf{x}\|_{t-1}^2.
\end{aligned} \tag{40}$$

We will assume that for any $t \geq 2$,

$$\|\mathbf{x}_t - \mathbf{x}_t^*\|_t \leq \frac{1}{6} \tag{41}$$

which will be proved at the end of the analysis.

According to the Taylor's theorem, for any $t \geq 2$, we have

$$f_t(\mathbf{x}_t) - f_t(\mathbf{x}_t^*) = \frac{1}{2} (\mathbf{x}_t - \mathbf{x}_t^*)^\top \nabla^2 f_t(\xi_t) (\mathbf{x}_t - \mathbf{x}_t^*)$$

where ξ_t is a point on the line segment between \mathbf{x}_t and \mathbf{x}_t^* . Now, using the property of self-concordant functions, we have

$$\nabla^2 f_t(\xi_t) = \nabla^2 f_t(\mathbf{x}_t^* + \xi_t - \mathbf{x}_t^*) \stackrel{(37)}{\preceq} \frac{1}{(1 - \|\xi_t - \mathbf{x}_t^*\|_t)^2} \nabla^2 f_t(\mathbf{x}_t^*) \preceq \frac{1}{(1 - \|\mathbf{x}_t - \mathbf{x}_t^*\|_t)^2} \nabla^2 f_t(\mathbf{x}_t^*)$$

where we use the inequality in (41) to ensure $\|\mathbf{x}_t - \mathbf{x}_t^*\|_t \leq 1$. We thus have

$$f_t(\mathbf{x}_t) - f_t(\mathbf{x}_t^*) \leq \frac{\|\mathbf{x}_t - \mathbf{x}_t^*\|_t^2}{2(1 - \|\mathbf{x}_t - \mathbf{x}_t^*\|_t)^2} \stackrel{(41)}{\leq} \|\mathbf{x}_t - \mathbf{x}_t^*\|_t^2.$$

As a result

$$\sum_{t=1}^T f_t(\mathbf{x}_t) - f_t(\mathbf{x}_t^*) \leq f_1(\mathbf{x}_1) - f_1(\mathbf{x}_1^*) + \sum_{t=2}^T \|\mathbf{x}_t - \mathbf{x}_t^*\|_t^2. \tag{42}$$

We first bound the dynamic regret by \mathcal{S}_T^* . To this end, we have

$$\sum_{t=2}^T \|\mathbf{x}_t - \mathbf{x}_t^*\|_t^2 \leq \sum_{t=2}^T 2 (\|\mathbf{x}_t - \mathbf{x}_{t-1}^*\|_t^2 + \|\mathbf{x}_t^* - \mathbf{x}_{t-1}^*\|_t^2) \stackrel{(40)}{\leq} 2\mu \sum_{t=2}^T \|\mathbf{x}_t - \mathbf{x}_{t-1}^*\|_{t-1}^2 + 2\mathcal{S}_T^*. \tag{43}$$

We proceed to bound $\sum_{t=2}^T \|\mathbf{x}_t - \mathbf{x}_{t-1}^*\|_{t-1}^2$. Since \mathbf{x}_t is derived by applying the damped Newton method multiple times to the initial solution \mathbf{x}_{t-1} , we need to first bound $\lambda_{t-1}(\mathbf{x}_{t-1})$. To this end, we establish the following lemma.

Lemma 5. *Let $f(\mathbf{x})$ be a self-concordant function, and $\mathbf{x}^* = \operatorname{argmin}_{\mathbf{x}} f(\mathbf{x})$. If $\|\mathbf{u} - \mathbf{x}^*\|_{\mathbf{x}^*} < 1/2$, we have*

$$\lambda(\mathbf{u}) \leq \frac{1}{1 - 2\|\mathbf{u} - \mathbf{x}^*\|_{\mathbf{x}^*}} \|\mathbf{u} - \mathbf{x}^*\|_{\mathbf{x}^*}.$$

The above lemma implies

$$\lambda_{t-1}(\mathbf{x}_{t-1}) \leq \frac{1}{1 - 2\|\mathbf{x}_{t-1} - \mathbf{x}_{t-1}^*\|_{t-1}} \|\mathbf{x}_{t-1} - \mathbf{x}_{t-1}^*\|_{t-1} \stackrel{(41)}{\leq} \min\left(\frac{3}{2}\|\mathbf{x}_{t-1} - \mathbf{x}_{t-1}^*\|_{t-1}, \frac{1}{4}\right). \tag{44}$$

Recall the updating rule

$$\mathbf{z}_{t-1}^{j+1} = \mathbf{z}_{t-1}^j - \frac{1}{1 + \lambda_{t-1}(\mathbf{z}_{t-1}^j)} \left[\nabla^2 f_{t-1}(\mathbf{z}_{t-1}^j) \right]^{-1} \nabla f_{t-1}(\mathbf{z}_{t-1}^j), \quad j = 1, \dots, K.$$

From Lemma 4, we have

$$\lambda_{t-1}(\mathbf{z}_{t-1}^{j+1}) \stackrel{(39)}{\leq} 2\lambda_{t-1}^2(\mathbf{z}_{t-1}^j), \quad j = 1, \dots, K.$$

Since $\lambda_{t-1}(\mathbf{z}_{t-1}^1) = \lambda_{t-1}(\mathbf{x}_{t-1}) \leq 1/4$. By induction, it is easy to verify

$$\lambda_{t-1}(\mathbf{z}_{t-1}^j) \leq \frac{1}{4}, \quad j = 1, \dots, K, K+1. \quad (45)$$

Therefore,

$$\lambda_{t-1}(\mathbf{x}_t) = \lambda_{t-1}(\mathbf{z}_{t-1}^{K+1}) \leq \frac{1}{2} \lambda_{t-1}(\mathbf{z}_{t-1}^K) \leq \dots \leq \frac{1}{2^K} \lambda_{t-1}(\mathbf{z}_{t-1}^1) = \frac{1}{2^K} \lambda_{t-1}(\mathbf{x}_{t-1}). \quad (46)$$

Again, using Lemma 4, we have

$$\|\mathbf{x}_t - \mathbf{x}_{t-1}^*\|_{t-1} \stackrel{(38)}{\leq} \frac{\lambda_{t-1}(\mathbf{x}_t)}{1 - \lambda_{t-1}(\mathbf{x}_t)} \stackrel{(45),(46)}{\leq} \frac{4}{3} \frac{1}{2^K} \lambda_{t-1}(\mathbf{x}_{t-1}) \stackrel{(44)}{\leq} \frac{2}{2^K} \|\mathbf{x}_{t-1} - \mathbf{x}_{t-1}^*\|_{t-1}$$

implying

$$\|\mathbf{x}_t - \mathbf{x}_{t-1}^*\|_{t-1}^2 \leq \frac{4}{4^K} \|\mathbf{x}_{t-1} - \mathbf{x}_{t-1}^*\|_{t-1}^2. \quad (47)$$

Combining (43) with (47), we have

$$\begin{aligned} \sum_{t=2}^T \|\mathbf{x}_t - \mathbf{x}_t^*\|^2 &\leq \frac{8\mu}{4^K} \sum_{t=3}^T \|\mathbf{x}_{t-1} - \mathbf{x}_{t-1}^*\|_{t-1}^2 + 2\mu \|\mathbf{x}_2 - \mathbf{x}_1^*\|_1^2 + 2\mathcal{S}_T^* \\ &\leq \frac{1}{2} \sum_{t=2}^T \|\mathbf{x}_t - \mathbf{x}_t^*\|^2 + 2\mu \|\mathbf{x}_2 - \mathbf{x}_1^*\|_1^2 + 2\mathcal{S}_T^* \end{aligned} \quad (48)$$

where we use the fact $\frac{8\mu}{4^K} \leq 1/2$. From (48), we have

$$\sum_{t=2}^T \|\mathbf{x}_t - \mathbf{x}_t^*\|^2 \leq 4\mu \|\mathbf{x}_2 - \mathbf{x}_1^*\|_1^2 + 4\mathcal{S}_T^* \stackrel{(12)}{\leq} \frac{1}{36} + 4\mathcal{S}_T^*. \quad (49)$$

Substituting (49) into (42), we obtain

$$\sum_{t=1}^T f_t(\mathbf{x}_t) - f_t(\mathbf{x}_t^*) \leq 4\mathcal{S}_T^* + f_1(\mathbf{x}_1) - f_1(\mathbf{x}_1^*) + \frac{1}{36}.$$

Next, we bound the dynamic regret by \mathcal{P}_T^* . From (41) and (42), we immediately have

$$\sum_{t=1}^T f_t(\mathbf{x}_t) - f_t(\mathbf{x}_t^*) \leq f_1(\mathbf{x}_1) - f_1(\mathbf{x}_1^*) + \frac{1}{6} \sum_{t=2}^T \|\mathbf{x}_t - \mathbf{x}_t^*\|_t. \quad (50)$$

To bound the last term, we have

$$\begin{aligned} \sum_{t=2}^T \|\mathbf{x}_t - \mathbf{x}_t^*\|_t &\leq \sum_{t=2}^T (\|\mathbf{x}_t - \mathbf{x}_{t-1}^*\|_t + \|\mathbf{x}_t^* - \mathbf{x}_{t-1}^*\|_t) \\ &\stackrel{(40)}{\leq} \sqrt{\mu} \sum_{t=3}^T \|\mathbf{x}_t - \mathbf{x}_{t-1}^*\|_{t-1} + \sqrt{\mu} \|\mathbf{x}_2 - \mathbf{x}_1^*\|_1 + \mathcal{P}_T^* \\ &\stackrel{(47),(12)}{\leq} \sqrt{\frac{4\mu}{4^K}} \sum_{t=3}^T \|\mathbf{x}_{t-1} - \mathbf{x}_{t-1}^*\|_{t-1} + \frac{1}{12} + \mathcal{P}_T^* \\ &\leq \frac{1}{2} \sum_{t=2}^T \|\mathbf{x}_t - \mathbf{x}_t^*\|_t + \frac{1}{12} + \mathcal{P}_T^* \end{aligned}$$

which implies

$$\sum_{t=2}^T \|\mathbf{x}_t - \mathbf{x}_t^*\|_t \leq \frac{1}{6} + 2\mathcal{P}_T^*. \quad (51)$$

Combining (50) and (51), we have

$$\sum_{t=1}^T f_t(\mathbf{x}_t) - f_t(\mathbf{x}_t^*) \leq \frac{1}{3} \mathcal{P}_T^* + f_1(\mathbf{x}_1) - f_1(\mathbf{x}_1^*) + \frac{1}{36}.$$

Finally, we prove that the inequality in (41) holds. For $t = 2$, we have

$$\|\mathbf{x}_2 - \mathbf{x}_2^*\|_2^2 \leq 2\|\mathbf{x}_2 - \mathbf{x}_1^*\|_2^2 + 2\|\mathbf{x}_1^* - \mathbf{x}_2^*\|_2^2 \stackrel{(11),(40)}{\leq} 2\mu\|\mathbf{x}_2 - \mathbf{x}_1^*\|_1^2 + \frac{1}{72} \stackrel{(12)}{\leq} \frac{1}{36}.$$

Now, we suppose (41) is true for $t = 2, \dots, k$. We show (41) holds for $t = k + 1$. We have

$$\begin{aligned} \|\mathbf{x}_{k+1} - \mathbf{x}_{k+1}^*\|_{k+1}^2 &\leq 2\|\mathbf{x}_{k+1} - \mathbf{x}_k^*\|_{k+1}^2 + 2\|\mathbf{x}_k^* - \mathbf{x}_{k+1}^*\|_{k+1}^2 \\ &\stackrel{(11),(40)}{\leq} 2\mu\|\mathbf{x}_{k+1} - \mathbf{x}_k^*\|_k^2 + \frac{1}{72} \stackrel{(47)}{\leq} \frac{8\mu}{4K}\|\mathbf{x}_k - \mathbf{x}_k^*\|_k^2 + \frac{1}{72} \leq \frac{1}{2}\|\mathbf{x}_k - \mathbf{x}_k^*\|_k^2 + \frac{1}{72} \leq \frac{1}{36}. \end{aligned}$$

I Proof of Lemma 5

By the mean value theorem for vector-valued functions, we have

$$\nabla f(\mathbf{u}) - \nabla f(\mathbf{x}^*) = \int_0^1 \nabla^2 f(\mathbf{x}^* + \tau(\mathbf{u} - \mathbf{x}^*)) (\mathbf{u} - \mathbf{x}^*) d\tau. \quad (52)$$

Define

$$g(\mathbf{x}) = \mathbf{x}^\top [\nabla^2 f(\mathbf{u})]^{-1} \mathbf{x}$$

which is a convex function of \mathbf{x} . Then, we have

$$\begin{aligned} \lambda^2(\mathbf{u}) &= \left\langle \nabla f(\mathbf{u}), [\nabla^2 f(\mathbf{u})]^{-1} \nabla f(\mathbf{u}) \right\rangle = g(\nabla f(\mathbf{u})) \\ &\stackrel{(52)}{\leq} g\left(\int_0^1 \nabla^2 f(\mathbf{x}^* + \tau(\mathbf{u} - \mathbf{x}^*)) (\mathbf{u} - \mathbf{x}^*) d\tau\right) \leq \int_0^1 g(\nabla^2 f(\mathbf{x}^* + \tau(\mathbf{u} - \mathbf{x}^*)) (\mathbf{u} - \mathbf{x}^*)) d\tau \end{aligned} \quad (53)$$

where the last step follows from Jensen's inequality.

Define $\xi_\tau = \mathbf{x}^* + \tau(\mathbf{u} - \mathbf{x}^*)$ which lies in the line segment between \mathbf{u} and \mathbf{x}^* . In the following, we will provide an upper bound for

$$g(\nabla^2 f(\xi_\tau)(\mathbf{u} - \mathbf{x}^*)) = (\mathbf{u} - \mathbf{x}^*)^\top \nabla^2 f(\xi_\tau) [\nabla^2 f(\mathbf{u})]^{-1} \nabla^2 f(\xi_\tau)(\mathbf{u} - \mathbf{x}^*).$$

Following Lemma 4, we have

$$\nabla^2 f(\xi_\tau) = \nabla^2 f(\mathbf{x}^* + \xi_\tau - \mathbf{x}^*) \stackrel{(37)}{\preceq} \frac{1}{(1 - \|\xi_\tau - \mathbf{x}^*\|_{\mathbf{x}^*})^2} \nabla^2 f(\mathbf{x}^*) \preceq \frac{1}{(1 - \|\mathbf{u} - \mathbf{x}^*\|_{\mathbf{x}^*})^2} \nabla^2 f(\mathbf{x}^*), \quad (54)$$

$$\|\mathbf{u} - \xi_\tau\|_{\xi_\tau}^2 \stackrel{(54)}{\leq} \frac{\|\mathbf{u} - \xi_\tau\|_{\mathbf{x}^*}^2}{(1 - \|\mathbf{u} - \mathbf{x}^*\|_{\mathbf{x}^*})^2} \leq \frac{\|\mathbf{u} - \mathbf{x}^*\|_{\mathbf{x}^*}^2}{(1 - \|\mathbf{u} - \mathbf{x}^*\|_{\mathbf{x}^*})^2} < 1, \quad (55)$$

$$\nabla^2 f(\mathbf{u}) = \nabla^2 f(\xi_\tau + \mathbf{u} - \xi_\tau) \stackrel{(37)}{\succeq} (1 - \|\mathbf{u} - \xi_\tau\|_{\xi_\tau})^2 \nabla^2 f(\xi_\tau) \stackrel{(55)}{\succeq} \left(\frac{1 - 2\|\mathbf{u} - \mathbf{x}^*\|_{\mathbf{x}^*}}{1 - \|\mathbf{u} - \mathbf{x}^*\|_{\mathbf{x}^*}}\right)^2 \nabla^2 f(\xi_\tau). \quad (56)$$

As a result

$$\begin{aligned} g(\nabla^2 f(\xi_\tau)(\mathbf{u} - \mathbf{x}^*)) &\stackrel{(56)}{\leq} \left(\frac{1 - \|\mathbf{u} - \mathbf{x}^*\|_{\mathbf{x}^*}}{1 - 2\|\mathbf{u} - \mathbf{x}^*\|_{\mathbf{x}^*}}\right)^2 \langle (\mathbf{u} - \mathbf{x}^*), \nabla^2 f(\xi_\tau)(\mathbf{u} - \mathbf{x}^*) \rangle \\ &\stackrel{(54)}{\leq} \frac{1}{(1 - 2\|\mathbf{u} - \mathbf{x}^*\|_{\mathbf{x}^*})^2} \|\mathbf{u} - \mathbf{x}^*\|_{\mathbf{x}^*}^2. \end{aligned} \quad (57)$$

We complete the proof by substituting (57) into (53).